

Geometric crystals and set-theoretical solutions to the quantum Yang-Baxter equation

Pavel Etingof

Massachusetts Institute of Technology

Department of Mathematics, Rm 2-176

Cambridge, MA 02139

email: etingof@math.mit.edu

February 1, 2008

1 Introduction

The notion of a geometric crystal was introduced and developed recently in [2], motivated by the needs of representation theory of p -adic groups. It is shown in [1, 2] that some particular geometric crystals give rise to interesting set-theoretical solutions R of the quantum Yang-Baxter equation, which satisfy the involutivity (or unitarity) condition $R^{21}R = 1$ (more precisely, they give rise to “rational” set theoretical solutions, see below).

On the other hand, involutive set-theoretical solutions of the quantum Yang-Baxter equation are studied in [4], following a suggestion of Drinfeld, [3]. It turned out in [4] that there is an especially nice theory of such solutions if an additional nondegeneracy condition is satisfied. In particular, in this case one can define the structure group G_R of a solution (X, R) , which acts on X . The image G_R^0 of G_R in $\text{Aut}(X)$ is called the reduced structure group. The complexity of this group, in a sense, characterizes the complexity of the solution (X, R) .

In this note we show that the maps (X, R) arising from the geometric crystals of [1],[2] are nondegenerate, and give a new proof that they satisfy the quantum Yang-Baxter equation and the involutivity condition. Then we calculate the reduced structure group of (X, R) and show that it is a subgroup of the group $PGL_n(\mathbf{C}(\lambda))$. We also give a new, direct proof of Theorem 8.9 of [1].

Remark. There is a seemingly technical, but in fact fundamental point that needs to be stressed. The nondegenerate solutions coming from geometric crystals live *not* in the category of sets (like the solutions from [4]) but in the category of irreducible algebraic varieties, where morphisms are *birational* maps. In particular, they *cannot* be viewed as usual

nondegenerate set-theoretical solutions, studied in [4]; they are similar but yet essentially new, more complicated objects. As a result, the theory of [4] has to be generalized to the “rational” case, to be applicable to geometric crystals. This generalization is not straightforward, and it is not entirely clear which parts of the theory survive and which do not. This seems to be an interesting problem for future research.

Acknowledgments. The author thanks A.Braverman and D.Kazhdan for useful discussions. This work was partially supported by the NSF grant DMS-9988796, and conducted in part for the Clay Mathematics Institute.

2 Rational set-theoretical R-matrices

Let us generalize the setting of [4] to the “rational” case.

Let X be an irreducible algebraic variety over $\mathbf{C}$. Let R be a birational isomorphism of $X \times X$ to itself.

Let us write $R(x, y)$, $x, y \in X$, as

$$R(x, y) = (f_y(x), g_x(y)). \quad (1)$$

The rational mappings $f_z, g_z : X \rightarrow X$ are well defined for generic $z \in X$.

Definition 2.1 *R is said to be nondegenerate if f_z, g_z are defined and are birational isomorphisms for all z .*

For example, $R = 1$ is nondegenerate but $R = P$ (the permutation) is not.

Definition 2.2 *R is said to be involutive if $R^{21}R = 1$.*

The quantum Yang-Baxter equation for R is the equation

$$R_{12}R_{13}R_{23} = R_{23}R_{13}R_{12} \quad (2)$$

between birational isomorphisms of X^3 .

Definition 2.3 *A solution R of the quantum Yang-Baxter equation will be called a rational set-theoretical R-matrix.*

For an irreducible variety Y , let $\text{Bir}(Y)$ be the group of birational automorphisms of Y . An involutive R -matrix R defines, for each N , a homomorphism $\rho_N^R : S_N \rightarrow \text{Bir}(X^N)$, given by $\rho((i, i+1)) = PR_{i, i+1}$. For example, for $R = 1$ this is the usual action by permutations, and for $R = P$ the trivial action.

Proposition 2.4 *If R is nondgenerate then ρ_N^R is conjugate to the usual action ρ_N^1 . Namely, $\rho_N^R = J_N^{-1} \rho_N^1 J_N$, where $J_N \in \text{Bir}(X^N)$ is given by*

$$J_N(x_1, \dots, x_N) = (f_{x_N} \dots f_{x_2}(x_1), \dots, f_{x_N}(x_{N-1}), x_N). \quad (3)$$

Proof: As in [4], Section 1.

Let R be an involutive nondgenerate rational set-theoretical R-matrix. Let $U \subset X^2$ be the domain of definition of R , i.e. the largest open set such that R is regular on U . Let G_R be the group generated by the points of X with defining relations

$$xy = y'x' \quad (4)$$

if $(x, y) \in U$ and $R(x, y) = (x', y')$. The group G_R is called the structure group of (X, R) .

Proposition 2.5 *The group G_R acts on X by birational transformations in two ways: $z \rightarrow f_z^{-1}$ and $z \rightarrow g_z$.*

Proof: As in [4], Section 2.

Let us denote the image in $\text{Bir}(X)$ of the first action by G_R^+ and of the second one by G_R^- . These groups will be called reduced structure groups of (X, R) .

3 Rational set-theoretical R-matrices arising from geometric crystals.

Let us recall the construction of [1, 2] (see [1], Section 8).

Let $\mathcal{T} = (\mathbf{C}^*)^n$. Elements of $\mathcal{T}$ will be written as $\mathbf{t} = [t_1, \dots, t_n]$, $t_i \in \mathbf{C}^*$.

For $k \geq 2$ let

$$\Delta_k(\mathbf{x}, \mathbf{y}) = x_1 \dots x_{k-1} + x_1 \dots x_{k-2} y_k + \dots + y_2 \dots y_k. \quad (5)$$

(we agree that $\Delta_1 = 1$). Let

$$\eta(\mathbf{x}, \mathbf{y}) = \frac{x_1 \dots x_n - y_1 \dots y_n}{\Delta_n(\mathbf{x}, \mathbf{y})} \quad (6)$$

The following proposition is easy.

Proposition 3.1 *[1, 2] There exists a unique rational map $R : \mathcal{T}^2 \rightarrow \mathcal{T}^2$,*

$$R(\mathbf{x}, \mathbf{y}) = (\mathbf{x}', \mathbf{y}'), \quad (7)$$

such that

$$x'_1 \dots x'_k = y_1 \dots y_k + \Delta_k(\mathbf{x}, \mathbf{y}) \eta(\mathbf{x}, \mathbf{y}), \quad (8)$$

and

$$x'_i y'_i = x_i y_i, \quad (9)$$

$$\prod_{i=1}^n x_i = \prod_{i=1}^n x'_i. \quad (10)$$

Our first result is

Proposition 3.2 *R is nondegenerate.*

This proposition is proved in the next section.

Moreover, one has the following result, which is a part of Theorem 8.9 of [1], proved in [2] using geometric crystals.

Theorem 3.3 *R is an involutive rational set-theoretical R-matrix.*

We give a direct proof of this theorem in Section 5. We note that such a proof was known to the authors of [1], but as far as we know, it is unpublished.

Corollary 3.4 *The S_N action on $\mathcal{T}^N$ defined by R is conjugate to the usual one.*

Remark. Since in the construction of R one does not use subtraction (see [2]), one can regard R as a usual (i.e. not “rational”) set-theoretical R-matrix defined on the set of points of $\mathcal{T}$ with positive real coordinates. Another (essentially, equivalent) way to turn R into a usual set-theoretical R-matrix is to use tropicalization (see [2] and references therein). However, it is easy to see that these usual R-matrices will not be nondegenerate. We don’t expect that it is possible to interpret R as a usual (not “rational”) *nondegenerate* set-theoretical R-matrix, and believe that it is essential to use the “rational” generalization to attain nondegeneracy.

Our second result is

Proposition 3.5 *The groups G_R^+ , G_R^- are subgroups of $PGL_n(\mathbf{C}(\lambda))$. These subgroups are isomorphic to each other.*

This proposition is proved in Section 5. In the proof, we actually describe the groups G_R^+, G_R^- explicitly.

Remark. We want to stress that we don't understand the meaning of the groups $G_R^\pm$ and of Proposition 3.5 in the context of geometric crystals. We also don't know how to generalize the main construction of [4] (the bijective cocycle construction) to the rational case, so that it would produce the solutions considered here. It seems that such a generalization would be interesting and useful.

Finally, we give a direct proof of Theorem 8.9 of [1] (according to [1], a proof of this result without the theory of geometric crystals was previously unavailable). More specifically, we prove the following statement, which is what Theorem 8.9 of [1] claims in addition to Theorem 3.3.

Let $M = (\mathbf{C}^*)^{mn}$ be the set of n by m matrices with entries from $\mathbf{C}^*$. Using the R-matrix R defined in Section 3, we can define two symmetric group actions on M : one of S_m (on columns) and another of S_n (on rows).

Theorem 3.6 *The actions of S_n and S_m on M commute with each other.*

This theorem is proved in Section 6.

4 Proof of Proposition 3.2

Let us introduce functions on $T_i : \mathcal{T} \rightarrow \mathbf{C}^*$ by $T_0(\mathbf{t}) = 1$, and $T_i = T_{i-1}t_i \bmod n$, where $i \bmod n$ takes values $1, \dots, n$. In this way, $T_j(\mathbf{t})$ is defined for all integer j .

We have $\mathcal{T} = \cup_{c \in \mathbf{C}^*} \mathcal{T}_c$, where $\mathcal{T}_c = \{\mathbf{t} \in \mathcal{T} | T_n = c\}$. It is clear that for (almost) all c , the map $f_{\mathbf{z}}, g_{\mathbf{z}}$ map $\mathcal{T}_c$ to $\mathcal{T}_c$.

Define two compactifications of $\mathcal{T}_c$, the projective spaces P_+ and P_- . To do this, it is sufficient to define two open embeddings $j_+, j_- : \mathcal{T}_c \rightarrow \mathbf{C}P^{n-1}$. We define them by

$$j_+(\mathbf{t}) = (1, T_1(\mathbf{t}), \dots, T_{n-1}(\mathbf{t})); \quad j_-(\mathbf{t}) = (T_1(\mathbf{t})^{-1}, \dots, T_{n-1}(\mathbf{t})^{-1}, T_n(\mathbf{t})^{-1} = c^{-1}); \quad (11)$$

The projective coordinates on P_+, P_- will be denoted by $\mathbf{Z} = (Z_1, \dots, Z_n)$.

Now let us take $\mathbf{x} \in \mathcal{T}_a$, $\mathbf{y} \in \mathcal{T}_b$, and calculate $f_{\mathbf{y}} : \mathcal{T}_a \rightarrow \mathcal{T}_a$, $g_{\mathbf{x}} : \mathcal{T}_b \rightarrow \mathcal{T}_b$. Let $X_i = T_i(\mathbf{x}), Y_i = T_i(\mathbf{y})$.

Proposition 4.1 (i) *The map $f_{\mathbf{y}} : \mathcal{T}_a \rightarrow \mathcal{T}_a$ extends to a projective transformation of P_+ given by*

$$\mathbf{Z} \rightarrow A_f(\mathbf{y}, a)\mathbf{Z}, \quad (12)$$

where $\mathbf{Z}$ is understood as a column vector, and A_f is the matrix given by

$$A_f(\mathbf{y}, a)_{ij} = Y_{i-1}Y_j^{-1}a, i > j; \quad A_f(\mathbf{y}, a)_{ij} = Y_{i-1}Y_j^{-1}b, i \leq j, \quad (13)$$

for $1 \leq i, j \leq n$.

(ii) The map $g_{\mathbf{x}} : \mathcal{T}_b \rightarrow \mathcal{T}_b$ extends to a projective transformation of P_- given by

$$\mathbf{Z} \rightarrow A_g(\mathbf{x}, b)\mathbf{Z}, \quad (14)$$

where A_g is the matrix given by

$$A_g(\mathbf{x}, b)_{ij} = X_{j-1}X_i^{-1}a, i \geq j; \quad A_g(\mathbf{x}, b)_{ij} = X_{j-1}X_i^{-1}b, i < j, \quad (15)$$

for $1 \leq i, j \leq n$.

Proof: The proof is by a straightforward calculation.

Now it is easy to prove Proposition 3.2. For this, we need to check the invertibility of the matrices A_f and A_g . To prove the invertibility of this first matrix, it is sufficient to observe that $\det A_f(\mathbf{y}, a) = (b - a)^{n-1}$ (this is easily established by induction). The second matrix is the transpose of the first one, so the result follows.

5 Proof of Theorem 3.3 and Proposition 3.5

We start with proving Theorem 3.3. Let us calculate the matrix $A_f^{-1}(\mathbf{y}, a)$, computing the minors of A_f of size $n - 1$. We find that $(A_f^{-1})_{ij} = 0$ unless $j - i$ is 0 or 1 modulo n , and

$$(A_f^{-1})_{ii} = (b - a)^{-1}Y_iY_{i-1}^{-1}, \quad (A_f^{-1})_{i,i+1} = -(b - a)^{-1}, \quad (A_f^{-1})_{n,1} = -a(b - a)^{-1}. \quad (16)$$

The matrix A_g^{-1} is, as we mentioned, the transpose of A_f^{-1} .

This leads to universal expressions for f^{-1} and g^{-1} which do not depend of a, b and even of n , and have a “local” form (in terms of the indices involved). These expressions will be useful below, but also seem interesting by themselves.

Namely, let $\mathbf{X}, \mathbf{Y}$ denote infinite sequences consisting of all $X_i = T_i(\mathbf{x}), Y_i = T_i(\mathbf{y}), i \in \mathbb{Z}$. Consider the functions from pairs of sequences to sequences, given by the formulas

$$\phi_{\mathbf{Y}}(\mathbf{X})_i = Y_{i+1}Y_i^{-1}X_i - X_{i+1}, \quad \gamma_{\mathbf{X}}(\mathbf{Y})_i = (X_iX_{i-1}Y_i^{-1} - Y_{i-1}^{-1})^{-1}. \quad (17)$$

The formulas for A_f^{-1} and A_g^{-1} imply the following.

Proposition 5.1 *In terms of the projective coordinates X_i, Y_i , the maps $(\mathbf{x}, \mathbf{y}) \rightarrow f_{\mathbf{y}}^{-1}(\mathbf{x})$ and $(\mathbf{x}, \mathbf{y}) \rightarrow g_{\mathbf{x}}^{-1}(\mathbf{y})$ are given by the functions ϕ and γ respectively.*

Let us now use this proposition to show that R is involutive.

We must check that $\mathbf{y} = g_{f_{\mathbf{y}}^{-1}(\mathbf{x})}^{-1} \circ f_{\mathbf{x}}^{-1}(\mathbf{y})$, $\mathbf{x}, \mathbf{y} \in (\mathbf{C}^*)^n$.

According to Proposition 5.1, this reduces to the identity

$$(Y_{i+1}Y_i^{-1}X_i - X_{i+1})(Y_iY_{i-1}^{-1}X_{i-1} - X_i)^{-1}(X_{i+1}X_i^{-1}Y_i - Y_{i+1})^{-1} - (X_iX_{i-1}^{-1}Y_{i-1} - Y_i)^{-1} = Y_i^{-1}, \quad (18)$$

which is straightforward.

Now let us prove the quantum Yang-Baxter equation for R . It has three components. Let us prove the identity in the first components. It has the form

$$f_{\mathbf{y}}f_{\mathbf{x}} = f_{f_{\mathbf{y}}(\mathbf{x})}f_{g_{\mathbf{x}}(\mathbf{y})}. \quad (19)$$

After changes of variables and inversions (using the involutivity of R) we see that this identity is equivalent to the claim that the map $f_{f_{\mathbf{y}}^{-1}(\mathbf{x})}^{-1}f_{\mathbf{y}}^{-1}$ is a symmetric function of $\mathbf{x}, \mathbf{y}$. Applying this map to $\mathbf{z}$ using Proposition 5.1, we find that this map is given by the operator on sequences defined by

$$(B\mathbf{Z})_i = (Y_{i+2}Y_{i+1}^{-1}X_{i+1} - X_{i+2})(Y_{i+1}Y_i^{-1}X_i - X_{i+1})^{-1}(Y_{i+1}Y_iZ_i - Z_{i+1}) - (Y_{i+2}Y_{i+1}^{-1}Z_{i+1} - Z_{i+2}). \quad (20)$$

But the symmetry of this operator in $\mathbf{X}, \mathbf{Y}$ is clear: The coefficient of Z_i is $\frac{Y_{i+2}X_{i+1} - Y_{i+1}X_{i+2}}{Y_{i+1}X_i - X_{i+1}Y_i}$, while the coefficient of Z_{i+1} is

$$- (Y_{i+2}Y_{i+1}^{-1}X_{i+1} - X_{i+2})(Y_{i+1}Y_i^{-1}X_i - X_{i+1})^{-1} - Y_{i+2}Y_{i+1}^{-1}, \quad (21)$$

which equals $-\frac{Y_{i+2}X_i - Y_iX_{i+2}}{Y_{i+1}X_i - X_{i+1}Y_i}$. This proves the first component of QYBE.

The third component of QYBE is completely analogous to the first one. The second component follows automatically from the involutivity and the first component, as shown in Proposition 2.2c in [4] (the straightforward proof of this from [4] applies verbatim in the case of rational maps). Theorem 3.3 is proved.

Now let us prove Proposition 3.5. Recall that by definition, the group G_R^+ is a subgroup of $\text{Bir}(\mathcal{T})$ generated by $f_{\mathbf{y}}^{-1}$, $\mathbf{y} \in \mathcal{T}$. In other words, G_R^+ is generated by the matrix functions $A_f(\mathbf{z}, \lambda)^{-1}$ of the parameter $\lambda \in \mathbf{C}^*$ for all $\mathbf{z} \in \mathcal{T}$. Thus, $G_R^+ \subset PGL_n(\mathbf{C}(\lambda))$. Similarly, $G_R^- \subset PGL_n(\mathbf{C}(\lambda))$ is generated by the matrix functions $A_g(\mathbf{z}, \lambda)$, $\mathbf{z} \in \mathcal{T}$.

Finally, it is easy to see that the groups G_R^+, G_R^- are isomorphic, since $A_f(\mathbf{z}, \lambda)^T = A_g(\mathbf{z}, \lambda)$.

Proposition 3.5 is proved.

Remark. It is shown in [4] that for usual nondegenerate involutive set-theoretical R-matrices, the groups G_R^+, G_R^- are not only isomorphic but also conjugate in $\text{Aut}(X)$ by a certain bijection T , which intertwines f_x^{-1} and g_x . In our situation, however, the map T is not well defined because of the “rational” character of the R-matrices. Therefore, the isomorphism of G_R^+, G_R^- has to be checked independently.

6 Proof of Theorem 3.6

Consider an n by m matrix (z_{kl}) ($n, m > 1$) with entries from $\mathbf{C}^*$. Fix $i : 1 \leq i \leq m - 1$, and $j : 1 \leq j \leq n - 1$. We want to show that the involution $\sigma = PR_{j,j+1}$ acting on the rows of this matrix, commutes with the involution $\tau = PR_{i,i+1}$ acting on the columns. For this purpose, we will calculate the composition $\tau\sigma$, and show that it is “symmetric under transposing the matrix”, which would imply the commutativity.

The elements which are subject to change under the composition $\tau\sigma$ are only those in rows $j, j + 1$ or columns $i, i + 1$. Therefore, it makes sense to introduce separate notations for these elements: $x_k = z_{jk}, y_k = z_{j+1,k}, p_l = z_{li}, q_l = z_{l+1,i}$. Introduce X_k, Y_k as in Section 4, and similarly P_l, Q_l . The same entries after application of σ , respectively $\tau\sigma$, will be denoted by the same letters with $*$, respectively $**$. Let $X_m = a, Y_m = b, P_n = c, Q_n = d$. The running subscript k will be between 0 and m , and l between 0 and n . Since our rows and columns have four elements in common, we have the following four equations:

$$\frac{P_j}{P_{j-1}} = \frac{X_i}{X_{i-1}}, \frac{Q_j}{Q_{j-1}} = \frac{X_{i+1}}{X_i}, \frac{Q_{j+1}}{Q_j} = \frac{Y_{i+1}}{Y_i}, \frac{P_{j+1}}{P_j} = \frac{Y_i}{Y_{i-1}}. \quad (22)$$

Let us also introduce the notation

$$G_k(\mathbf{X}, \mathbf{Y}) = a \sum_{r=1}^k Y_r^{-1} X_{r-1} + b \sum_{r=k+1}^m Y_r^{-1} X_{r-1}, \quad (23)$$

and similarly

$$H_l(\mathbf{P}, \mathbf{Q}) = c \sum_{r=1}^l Q_r^{-1} P_{r-1} + d \sum_{r=l+1}^n Q_r^{-1} P_{r-1}. \quad (24)$$

According to section 4, we have

$$X_k^* = X_k \frac{G_0(\mathbf{X}, \mathbf{Y})}{G_k(\mathbf{X}, \mathbf{Y})}, \quad (25)$$

and

$$Y_k^* = Y_k \frac{G_k(\mathbf{X}, \mathbf{Y})}{G_0(\mathbf{X}, \mathbf{Y})}, \quad (26)$$

We also obviously have $P_l^* = P_l, Q_l^* = Q_l$ for $l \neq j$. Finally, for $l = j$, using formulas 22, it is easy to find that

$$P_j^* = P_j \frac{G_{i-1}(\mathbf{X}, \mathbf{Y})}{G_i(\mathbf{X}, \mathbf{Y})}, \quad (27)$$

and

$$Q_j^* = Q_j \frac{G_i(\mathbf{X}, \mathbf{Y})}{G_{i+1}(\mathbf{X}, \mathbf{Y})}, \quad (28)$$

Now let us calculate the parameters after application of $\tau\sigma$. We clearly have

$$X_k^{**} = X_k^* = X_k \frac{G_0(\mathbf{X}, \mathbf{Y})}{G_k(\mathbf{X}, \mathbf{Y})}, k \neq i. \quad (29)$$

and

$$Y_k^{**} = Y_k^* = Y_k \frac{G_k(\mathbf{X}, \mathbf{Y})}{G_0(\mathbf{X}, \mathbf{Y})}, k \neq i \quad (30)$$

Now let us compute Q_l^{**} . We have

$$Q_l^{**} = Q_l^* \frac{H_l(\mathbf{P}^*, \mathbf{Q}^*)}{H_0(\mathbf{P}^*, \mathbf{Q}^*)}. \quad (31)$$

Let us express this via P_r, Q_r . As we know, $P_r^* = P_r$ and $Q_r^* = Q_r$, except for $r = j$. Thus, if $l \neq j$, all the stars in this expression can be removed, except in the combination $(Q_j^*)^{-1}P_{j-1} + Q_{j+1}^{-1}P_j^*$, which occurs both in the numerator and the denominator.

Now, using the formulas for P_j^*, Q_j^* , we have

$$(Q_j^*)^{-1}P_{j-1} = Q_j^{-1}P_{j-1} + (a-b) \frac{Q_j^{-1}P_{j-1}Y_{i+1}^{-1}X_i}{G_i(\mathbf{X}, \mathbf{Y})}, \quad (32)$$

and

$$Q_{j+1}^{-1}P_j^* = Q_{j+1}^{-1}P_j - (a-b) \frac{Q_{j+1}^{-1}P_jY_i^{-1}X_{i-1}}{G_i(\mathbf{X}, \mathbf{Y})}. \quad (33)$$

But it is easy to check using equations 22 that

$$Q_j^{-1}P_{j-1}Y_{i+1}^{-1}X_i = Q_{j+1}^{-1}P_jY_i^{-1}X_{i-1}. \quad (34)$$

This implies that

$$(Q_j^*)^{-1}P_{j-1} + Q_{j+1}^{-1}P_j^* = Q_j^{-1}P_{j-1} + Q_{j+1}^{-1}P_j, \quad (35)$$

and hence

$$Q_l^{**} = Q_l \frac{H_l(\mathbf{P}, \mathbf{Q})}{H_0(\mathbf{P}, \mathbf{Q})}. \quad (36)$$

We now notice the desired symmetry with the expression for Y_k^{**} , $k \neq i$, under the relabeling $(P, Q, l, j, n, c, d, H) \rightarrow (X, Y, k, i, m, a, b, G)$. Similarly,

$$P_l^{**} = P_l \frac{H_0(\mathbf{P}, \mathbf{Q})}{H_l(\mathbf{P}, \mathbf{Q})}, \quad (37)$$

which is symmetric to the expression for X_k^{**} .

It remains to compute $X_i^{**}, Y_i^{**}, P_j^{**}$ and Q_j^{**} and make sure the desired symmetry is present. Using the symmetric images of the equations for 27, 28 for P_j^*, Q_j^* , we get

$$X_i^{**} = X_i^* \frac{H_{j-1}(\mathbf{P}^*, \mathbf{Q}^*)}{H_j(\mathbf{P}^*, \mathbf{Q}^*)} = \quad (38)$$

$$X_i \frac{G_0(\mathbf{X}, \mathbf{Y})}{G_i(\mathbf{X}, \mathbf{Y})} \cdot \frac{H_{j-1}(\mathbf{P}, \mathbf{Q})}{H_j(\mathbf{P}^*, \mathbf{Q}^*)} \quad (39)$$

(the stars in the numerator are removed by formula 35). Now, by formulas 32, 33, we have

$$H_j(\mathbf{P}^*, \mathbf{Q}^*) = c \sum_{r=1}^j (Q_r^*)^{-1} P_{r-1}^* + d \sum_{r=j+1}^n (Q_r^*)^{-1} P_{r-1}^* = \frac{S}{G_i(\mathbf{X}, \mathbf{Y})}, \quad (40)$$

where

$$S = (c - d)(a - b)U + G_i(\mathbf{X}, \mathbf{Y})H_j(\mathbf{P}, \mathbf{Q}), \quad U = Q_j^{-1}P_{j-1}Y_{i+1}^{-1}X_i = Q_{j+1}^{-1}P_jY_i^{-1}X_{i-1}. \quad (41)$$

Thus,

$$X_i^{**} = X_i S^{-1} G_0(\mathbf{X}, \mathbf{Y}) H_{j-1}(\mathbf{P}, \mathbf{Q}). \quad (42)$$

The element P_j^{**} can now be found from equations 22:

$$P_j^{**} = X_i^{**} P_{j-1}^{**} (X_{i-1}^{**})^{-1} = P_j S^{-1} H_0(\mathbf{P}, \mathbf{Q}) G_{i-1}(\mathbf{X}, \mathbf{Y}). \quad (43)$$

Now it is apparent that X_i^{**} is symmetric to P_j^{**} under the symmetry $(P, Q, l, j, n, c, d, H) \rightarrow (X, Y, k, i, m, a, b, G)$. Namely, it follows from the fact that the expression U , and hence S , is invariant under this symmetry.

Finally, as we showed before, the expression $X_i^{**} Y_i^{**} = X_{i-1}^{**} Y_{i-1}^{**} P_{j+1}^{**} / P_{j-1}^{**}$ is symmetric to $P_{j-1}^{**} Q_{j-1}^{**} X_{i+1}^{**} / X_{i-1}^{**} = P_j^{**} Q_j^{**}$, which implies that Y_i^{**} is symmetric to Q_j^{**} . The theorem is proved.

References

- [1] A.Braverman and D.Kazhdan (with an appendix by V.Vologodsky), γ -functions of representations and lifting, math.AG 9912208, Geom. Funct. Anal. 2000, Special Volume, Part I, 237–278.
- [2] A.Berenstein and D.Kazhdan, Geometric and unipotent crystals, math 9912105, Geom. Funct. Anal. 2000, Special Volume, Part I, 188–236.
- [3] V.Drinfeld, On some unsolved problems in quantum group theory, in “Quantum groups” (Leningrad 1990), Lect. Notes in Math, 1510, Springer, Berlin, p. 1-8, 1992.
- [4] P.Etingof, T.Schedler, A.Soloviev, Set-theoretical solutions to the quantum Yang-Baxter equation, Duke Math.J. v.100, issue 2, 1999.